



Impact of Feeding Postbiotics and Paraprobiotics Produced From *Lactiplantibacillus plantarum* on Colon Mucosa Microbiota in Broiler Chickens

Yohanna Danladi¹, Teck Chwen Loh^{1,2*}, Hooi Ling Foo^{3,4*}, Henny Akit^{1,4}, Nur Aida Md Tamrin¹ and Azizi Mohammad Naeem¹

OPEN ACCESS

Edited by:

Francesco Gai,
Italian National Research Council, Italy

Reviewed by:

Claudia Cecilia Márquez Mota,
National Autonomous University of
Mexico, Mexico
Yan Zhao,
Wenzhou Academy of Agricultural
Sciences, China

*Correspondence:

Teck Chwen Loh
tcloh@upm.edu.my
Hooi Ling Foo
hlfoo@upm.edu.my

Specialty section:

This article was submitted to
Animal Nutrition and Metabolism,
a section of the journal
Frontiers in Veterinary Science

Received: 21 January 2022

Accepted: 14 February 2022

Published: 29 March 2022

Citation:

Danladi Y, Loh TC, Foo HL, Akit H, Md Tamrin NA and Mohammad Naeem A (2022) Impact of Feeding Postbiotics and Paraprobiotics Produced From *Lactiplantibacillus plantarum* on Colon Mucosa Microbiota in Broiler Chickens. *Front. Vet. Sci.* 9:859284. doi: 10.3389/fvets.2022.859284

¹ Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Malaysia, ² Institute of Tropical Agriculture and Food Security, Universiti Putra Malaysia, Serdang, Malaysia, ³ Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Malaysia, ⁴ Institute of Bioscience, Universiti Putra Malaysia, Serdang, Malaysia

This study was conducted to evaluate the impact of feeding postbiotics and paraprobiotics produced from *Lactiplantibacillus plantarum* on colon mucosa microbiota in broiler chickens. In this study, 336 one-day-old COBB 500 chicks were randomly allotted to eight treatment groups and replicated six times with seven birds per replicate. The treatment included T1 (Negative control) = Basal diet, T2 (Positive control) = Basal diet + 0.01% oxytetracycline, T3 = Basal diet + 0.2% postbiotic TL1, T4 = Basal diet + 0.2% postbiotic RS5, T5 = Basal diet + 0.2% paraprobiotic RG11, T6 = Basal diet + 0.2% postbiotic RI11, T7 = Basal diet + 0.2% paraprobiotic RG14, and T8 = Basal diet + 0.2% paraprobiotic RI11. There were reported changes in the bacterial community using 16S rRNA sequencing of the colon mucosa. The results of the sequencing of 16S rRNA genes in the colon mucosa samples indicated that compared to birds fed the negative control diet, birds fed paraprobiotic RI11 diets were recorded to have a lower relative abundance of Proteobacteria, while those fed the positive control were recorded to have a higher proportion of Firmicutes. Also, lower *Enterococcus* was reported in paraprobiotic RI11, while the most abundant genus was *Bacteroides* in postbiotic TL1. This study revealed that supplementation of postbiotics and paraprobiotics in the diets of broilers demonstrated positive effects on the microbiota by supporting the increase of beneficial microbes like the *Firmicutes* while decreasing harmful microbes like the *Proteobacteria*. Therefore, this study has provided knowledge on the modification of chicken mucosa microbiota through the feeding of postbiotics and paraprobiotics.

Keywords: postbiotics, paraprobiotics, colon mucosa, microbiota, broiler

INTRODUCTION

The gastrointestinal tract (GIT) of chicken hosts a great microbial community, and its integrity play an important role in nutrient absorption, development of immunity, and disease resistance (1). When changes occur in the GIT microbiome, feed efficiency, productivity, and health of the birds can be influenced (2–4). The different sections of the GIT of chickens are heavily populated with complex microbiome (bacteria, fungi, Archaea, protozoa, and virus) dominated by bacteria (5). The gut microbiota can create a protective barrier by attaching to the epithelial walls of enterocytes, thereby reducing the possibility of colonising pathogenic bacteria (6).

Primarily, the commensal microbiota can benefit the host in terms of competitive exclusion of pathogens or non-indigenous microbes, immune stimulation and programming, and contributions to the host nutrition (7). In addition, they can stimulate the development of the immune system, including mucus layer, epithelial monolayer, intestinal immune cells (e.g., cytotoxic and helper T cells, immunoglobulin producing cells, and phagocytic cells), and the lamina propria (7–9). The gut mucus layer increases mucin secretion and epithelial cell turnover through the influence of the commensal microbiota. This helps keep the GIT lubricated while preventing microorganisms from taking over the intestinal epithelial cells of the host (1).

The microbiota in the hindgut (caeca and colon) can produce energy and nutrients like vitamins, amino acids, and short-chain fatty acids (SCFA) from the undigested feed, which are later available to the host (2, 7). The bacteriostatic properties of SCFA can destroy foodborne pathogens such as *Salmonella* sp. (10). Previous reports have already established that the pH of the colon becomes lower as a result of SCFA production, which inhibits the conversion of bile to secondary bile products (11).

Diets greatly impact the intestinal microbiome of chickens. Different constituents of diet that escape digestion and absorption by the host are being utilised as a substrate for growth by the bacteria (12). The diversity and composition of the gut microbiome in poultry are greatly impacted by diets (13).

The composition of the mucosa attached microbiota is influenced by several host factors such as the expression of specific adhesion sites on the enterocyte membrane, secretion of secretory immunoglobulins, and mucus production rate (1). It is well established that diet can alter luminal and mucosa-attached microbiota to influence gut health (3). The use of postbiotics as a replacement for antibiotics in broiler chicken diets has been well documented (14–17). Postbiotic has been defined as any factors resulting from the metabolic activity of a probiotic or any released molecules capable of conferring beneficial effects to the host directly or indirectly (18). Postbiotics are also known as non-viable bacterial products or metabolic by-products from probiotic microorganisms with biological activity in the host (18, 19). In contrast, the term paraprobiotics was coined to indicate the use of inactivated probiotic cells (non-viable) or probiotic cell fractions to confer health benefits to the host (20). Hence, paraprobiotics are also known as “inactivated probiotics” or ghost probiotics (20). The benefits of postbiotics

include inhibiting pathogenic bacteria growth, leading to efficient nutrient utilisation and improvement in growth (21–23). On the other hand, the benefits provided to the host by paraprobiotics include modulation of the immune system, whereby the cell wall components may boost the immunological responses (24, 25).

Extensive studies with postbiotics effects on targeted caecum microbial population have been carried out (14, 17, 26, 27). However, the effect of postbiotics and paraprobiotics on the colon bacterial microbiota is yet unknown. Similarly, postbiotics and paraprobiotics affect the microbiota living in the outer mucosa layer, which differs from the caecal microbiome. Mucosa microbiota found within the outer mucous layer plays an important role in the intestine (28, 29). Therefore, the colon mucosal genome was studied using next-generation sequencing (16S rRNA). This study aimed to evaluate the impact of postbiotics and paraprobiotics on the colon mucosal microbiota in broiler chickens.

MATERIALS AND METHODS

Postbiotic and Paraprobiotic Preparations

The active culture of *Lactiplantibacillus plantarum* strains was washed once with sterile 0.85% (w/v) NaCl (Merck, Darmstadt, Germany) solution and adjusted to 10^9 CFU/ml to be used as a 10% (v/v) inoculum according to the method described by Mohamad et al. (30). Both postbiotics and paraprobiotics were prepared according to the method described by Ooi et al. (31) using de Man, Rogosa, and Sharpe (MRS) medium and incubated at 30°C for 24 h under anaerobic conditions. As for postbiotic preparation, cell-free supernatant was collected as postbiotics after centrifugation at $10,000 \times g$ for 15 min at 4°C. The cell suspension of *L. plantarum* strains was frozen for 7 days at -30°C to produce paraprobiotics.

Broiler Chicken and Management

A total of 336 day-old COBB 500 chicks (DOCs) were obtained from a commercial hatchery. The DOCs were randomly distributed to 8 dietary treatments managed in a closed house system. The house temperature was set at $33^\circ\text{C} \pm 1^\circ\text{C}$ on day 1 and was gradually reduced to about $25^\circ\text{C} \pm 1^\circ\text{C}$ by day 15. The average relative humidity ranged between 60 and 75%. Each treatment group was replicated six times with seven birds per replicate and was managed in a 120 × 120 cm (length × width) pen cage. The dietary treatment included T1 (Negative control) = Basal diet, T2 (Positive control) = Basal diet + 0.01% oxytetracycline, T3 = Basal diet + 0.2% TL1 postbiotic, T4 = Basal diet + 0.2% RS5 postbiotic, T5 = Basal diet + 0.2% RG11 paraprobiotic, T6 = Basal diet + 0.2% RI11 postbiotic, T7 = Basal diet + 0.2% RG11 paraprobiotic, and T8 = Basal diet + 0.2% RI11 paraprobiotic. The birds were vaccinated against Newcastle disease and infectious bronchitis disease (ND-IB) through eye drop at 7 and 21 days. The infectious bursal disease (IBD) vaccination was done on day 14 by eye drop. Water and feed were offered *ad libitum* until day 35. The starter and finisher diets (Tables 1, 2) were offered from days 0 to 21 and days 22 until 35 days of age, respectively. The experiment was undertaken based on the guidelines approved by the

TABLE 1 | Nutrient composition of starter diets (days 1–21).

Ingredients	Treatment diets							
	T1	T2	T3	T4	T5	T6	T7	T8
Corn	47.50	47.49	47.20	47.20	47.20	47.20	47.20	47.20
Soybean meal	40.10	40.10	40.20	40.20	40.20	40.20	40.20	40.20
Wheat pollard	1.50	1.50	1.60	1.60	1.60	1.60	1.60	1.60
CPO	6.00	6.00	5.90	5.90	5.90	5.90	5.90	5.90
L-Lysine	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
dl-Methionine	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Dicalcium phosphate	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50
Calcium carbonate	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45
Choline chloride	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Salt	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
Mineral mix	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Vitamin mix	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Antioxidant	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Toxin binder	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Antibiotics	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Postbiotic TL1	0.00	0.00	0.20	0.00	0.00	0.00	0.00	0.00
Postbiotic RS5	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.00
Paraprobiotic RG11	0.00	0.00	0.00	0.00	0.20	0.00	0.00	0.00
Postbiotic RI11	0.00	0.00	0.00	0.00	0.00	0.20	0.00	0.00
Paraprobiotic RG14	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00
Paraprobiotic RI11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.20
Total	100	100	100	100	100	100	100	100
Calculated analysis								
ME (kcal/kg)	3,215.70	3,215.70	3,201.40	3,201.40	3,201.40	3,201.40	3,201.40	3,201.40
Protein (%)	22.00	22.00	22.03	22.03	22.03	22.03	22.03	22.03
Fat (%)	7.99	7.99	7.88	7.88	7.88	7.88	7.88	7.88
Fibre (%)	4.11	4.11	4.12	4.12	4.12	4.12	4.12	4.12
Calcium (%)	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08
Total Phos (%)	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89
Avail. P (%)	0.48	0.48	0.48	0.48	0.48	0.48	0.48	0.48

Negative control = T1 (Basal diet), positive control = T2 (Basal diet + 0.01% (w/w) Oxytetracycline), T3 = Basal diet + 0.2% (v/w) postbiotic TL1, T4 = Basal diet + 0.2% (v/w) Postbiotic RS5, T5 = Basal diet + 0.2% (v/w) paraprobiotic RG11, T6 = Basal diet + 0.2% (v/w) postbiotic RI11, T7 = Basal diet + 0.2% (v/w) paraprobiotic RG14, T8 = Basal diet + 0.2% (v/w) paraprobiotic RI11. Dicalcium phosphate 18%; vitamin premix provided per kilogram of diet: vitamin A 35,000 MIU; vitamin D3 9,000 MIU; vitamin E 90,000 g; vitamin K3 6,000 g; vitamin B1 7,000 g; vitamin B2 22,000 g; vitamin B6 12,000 g; vitamin B12 0.070 g; biotin 300,000 mg; pantothenic acid 35,000 g; nicotinic acid, 120,000 g; folic acid 3,000 g; phytase 25,000,000 FTU. Mineral mix provided per kilogram of diet: Se 0.2 g, Cu 15 g, Fe 80 g, I 1 g, Mn 100 g, Na 1.5 g, Zn 80 g, K 4 g, and Co 0.25 g. Antioxidant contains butylated hydroxyanisole (BHA). Toxin binder contains natural hydrated sodium calcium aluminium silicates to reduce the exposure of feed to mycotoxins. Oxytetracycline (200 mg/kg, purity \geq 64.7%, Y.S.P. Industries (M) SDN BHD). The diets were formulated using FeedLIVE International software (Nonthaburi, Thailand).

Institutional Animal Care and Use Committee of the Universiti Putra Malaysia (IACUC) with reference no. UPM/IACUC/AUP-R098/2018, which ensures that the care and use of animals for scientific purposes are humane and ethical.

Mucosa Bacterial Metagenomic DNA Sample Preparation

At the end of day 35 of the experiment, six chickens were randomly selected from each treatment. The chickens were slaughtered, and the mucosal scrapings from the colon were collected, quickly frozen, and later stored at -80°C until the time for analysis.

Bacterial Genomic DNA Extraction

The bacterial genomic DNA (gDNA) was extracted from the colonic mucosa samples using the NucleoSpin® DNA stool kit (Macherey-Nagel, GmbH and Co. KG, Düren, Germany). Approximately 200 mg of frozen colonic mucosa samples was lysed in ST1 buffer, the lysate was filtered using the NucleoSpin® Inhibitor Removal column, and Buffer ST2 was added to precipitate contaminants. Buffer ST3 was added to adjust the binding conditions, and the NucleoSpin® DNA Stool column was used to bind the DNA. The NucleoSpin® DNA Stool column was washed in four steps using buffers ST3, ST4, and ST5. Buffer SE was used to elute the DNA after the washing steps. DNA

TABLE 2 | Nutrient composition of finisher diets (days 22–35).

Ingredients	Treatment diets							
	T1	T2	T3	T4	T5	T6	T7	T8
Corn	51.60	51.59	51.60	51.60	51.60	51.60	51.60	51.60
Soybean meal	33.50	33.50	33.50	33.50	33.50	33.50	33.50	33.50
Wheat pollard	4.80	4.80	4.60	4.60	4.60	4.60	4.60	4.60
CPO	5.20	5.20	5.20	5.20	5.20	5.20	5.20	5.20
L-Lysine	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
dl-Methionine	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Dicalcium phosphate	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50
Calcium carbonate	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45
Choline chloride	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Salt	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
Mineral mix	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Vitamin mix	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Antioxidant	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Toxin binder	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Antibiotics	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Postbiotic TL1	0.00	0.00	0.20	0.00	0.00	0.00	0.00	0.00
Postbiotic RS5	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.00
Paraprobiotic RG11	0.00	0.00	0.00	0.00	0.20	0.00	0.00	0.00
Postbiotic RI11	0.00	0.00	0.00	0.00	0.00	0.20	0.00	0.00
Paraprobiotic RG14	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00
Paraprobiotic RI11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.20
Total	100	100	100	100	100	100	100	100
Calculated analysis								
ME (kcal/kg)	3,180.83	3,180.83	3,176.68	3,176.68	3,176.68	3,176.68	3,176.68	3,176.68
Protein (%)	19.92	19.92	19.89	19.89	19.89	19.89	19.89	19.89
Fat (%)	7.29	7.29	7.29	7.29	7.29	7.29	7.29	7.29
Fibre (%)	4.01	4.01	3.99	3.99	3.99	3.99	3.99	3.99
Calcium (%)	1.06	1.06	1.06	1.06	1.06	1.06	1.06	1.06
Total Phos (%)	0.89	0.89	0.88	0.88	0.88	0.88	0.88	0.88
Avail. P (%)	0.48	0.48	0.48	0.48	0.48	0.48	0.48	0.48

Negative control = T1 (Basal diet), positive control = T2 [Basal diet + 0.01% (w/w) Oxytetracycline], T3 = Basal diet + 0.2% (v/w) postbiotic TL1, T4 = Basal diet + 0.2% (v/w) Postbiotic RS5, T5 = Basal diet + 0.2% (v/w) paraprobiotic RG11, T6 = Basal diet + 0.2% (v/w) postbiotic RI11, T7 = Basal diet + 0.2% (v/w) paraprobiotic RG14, and T8 = Basal diet + 0.2% (v/w) paraprobiotic RI11. Dicalcium phosphate 18%; vitamin premix provided per kilogram of diet: vitamin A 35,000 MIU; vitamin D3 9,000 MIU; vitamin E 90,000 g; vitamin K3 6,000 g; vitamin B1 7,000 g; vitamin B2 22,000 g; vitamin B6 12,000 g; vitamin B12 0.070 g; biotin 300,000 mg; pantothenic acid 35,000 g; nicotinic acid, 120,000 g; folic acid 3,000 g; phytase 25,000,000 FTU. Mineral mix provided per kilogram of diet: Se 0.2 g, Cu 15 g, Fe 80 g, I 1 g, Mn 100 g, Na 1.5 g, Zn 80 g, K 4 g, and Co 0.25 g. Antioxidant contains butylated hydroxyanisole (BHA). Toxin binder contains natural hydrated sodium calcium aluminium silicates to reduce the exposure of feed to mycotoxins. Oxytetracycline (200 mg/kg, purity \geq 64.7%, Y.S.P. Industries (M) SDN BHD). The diets were formulated using FeedLIVE International software (Nonthaburi, Thailand).

quality was verified using a NanoDrop 2000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA) with a concentration (260/280 nm ratio absorbance) of extracted DNA.

16S RRNA Sequencing of Colon Mucosa Microbiota Bacterial 16S V3–V4 Amplicon Sequencing

Twenty-four (24) purified gDNAs were sent to Apical Scientific Laboratory, Sdn Bhd, Seri Kembangan, Malaysia, for the sequencing. The quality of the purified DNAs was first monitored on 1% Tris–acetate–EDTA (TAE) agarose gel. The concentration of DNA was measured using a spectrophotometer (Implen

NanoPhotometer® N60/N50) and fluorometric quantification using iQuant™ Broad Range dsDNA Quantification Kit (Figure 1).

The purified gDNA that passed DNA sample QC was amplified using locus-specific sequence primers: 16S V3–V4 forward: CCTACGGGNGGCWGCAG, 16S V3–V4 reverse: GACTACHVGGGTATCTAATCC. All the PCRs were carried out with REDiant 2× PCR Master Mix (1st BASE). Library preparations were done in two stages. The first stage of the PCR of bacterial 16S rRNA gene of the selected regions (16S V3–V4) was amplified using locus-specific sequence primers with overhang adapters, as follows:




FIGURE 1 | Bacterial gDNA fluorometric quantification.

Forward overhang: 5' TCGTCGGCAGCGTCAGATGTGTAT AAGAGACAG-[locus-specific sequence]

Reverse overhang: 5' GTCTCGTGGGCTCGGAGATGTGTAT AAGAGACAG-[locus-specific sequence]. All the PCRs were done using KOD, Multi & Epi® (Toyobo, Osaka, Japan).

At the second stage of PCR, dual indices were attached to the amplicon PCR using Illumina Nextera XT Index Kit v2 following the manufacturer's protocols. The quality of the libraries was measured using Agilent Bioanalyzer 2100 System by Agilent DNA 1000 Kit and fluorometric quantification by Helixyte Green™ Quantifying Reagent.

The libraries were normalised and pooled according to the protocol recommended by Illumina and proceeded to sequence using the MiSeq platform using 300 PE.

A phylogenetic tree was created using a combination of Multiple Alignment using Fast Fourier Transform (MAFFT) and FastTree algorithms. The MAFFT algorithms were used to construct a multiple sequence alignment (MSA), which was then passed to FastTree to construct a phylogenetic tree based on maximum-likelihood nearest-neighbour interchanges (NNIs). In addition, FastTree utilises the CAT estimation. It uses heuristics to restrict the search for a better phylogenetic tree and estimates a rate of evolution for each site at lower memory consumption and faster inference times.

Data Analysis

The sequence adapters and low-quality reads were removed from the paired-end reads before the first 200,000 raw reads were

extracted using BBTools. Then, the forward and reverse reads were merged using QIIME. DADA2 pipeline (<https://benjjneb.github.io/dada2/>) was used to remove and correct error reads and to remove low-quality regions and chimeric errors. The resulting data were in the form of amplicon sequence variant (ASV) and was used in the next steps accordingly. The taxonomic classification was done using scikit-learn (<https://scikit-learn.org/stable/>) and naive Bayes classifier against database SILVA (release 132).

It is truly common in amplicon sequencing to involve a portion of the 16S rRNA gene or 18S gene, where the sequences are classified taxonomically. However, this involves few software and pipelines. The DADA2 pipeline comes with a naive Bayesian classifier that can classify large sequences across multiple ranks—from kingdom to genus—and provide an output in the form of taxonomy assignments with bootstrap confidence. It compares a set of taxonomically assigned sequences provided from formatted reference fasta files databases such as SILVA for ribosomal rRNA database and make individual taxonomic assignment (32). SILVA database (Release 132) was used to analyse the sequence similarity within the ASV reads with recommended parameters at a 97% similarity threshold (33).

Bioinformatic Analyses

Sequencing the region in 16S rRNA was done with a paired-end (PE) Illumina MiSeq platform that generates 300-bp raw reads. Sequence adapters and low-quality reads were removed from the raw reads using BBDuk (version 38.76). The raw reads are aligned

and merged using QIIME2 (version 2019.10). The Divisive Amplicon Denoising Algorithm 2 (DADA2) pipeline (version 1.14) was used to denoise as an attempt to remove and correct error reads and to remove low-quality regions and chimeric errors to obtain ASV (34). DADA2 pipeline was used in this analysis to substitute the traditional operational taxonomic unit (OTU) method. DADA2 method is more sensitive and specific and can detect real biological variation, which is usually missed by the OTU classification methods. DADA2 can accurately resolve sequence variants differing by just one nucleotide and present in as few as two reads, making this pipeline more precise, comprehensive, and reproducible (35).

Alpha diversity was measured through 5 indices—Observed, Chao1, Shannon, Simpson, and Fisher—to determine the richness and the diversity of the bacteria in the colon mucosa according to the different dietary treatments. A rarefaction curve plot of the number of species (species richness) was plotted as a function of the number of samples (sequence sample size).

Statistical Analysis

Statistical tests for sequencing analysis were done based on alpha diversity. Statistical tests were performed in R Studio version 3.6.2 by using the following packages:

- phyloseq (<https://www.bioconductor.org/packages/release/bioc/html/phyloseq.html>),
- vegan (<https://cran.r-project.org/web/packages/vegan/index.html>), and
- Venn Diagram (<https://cran.rproject.org/web/packages/VennDiagram/index.html>).

RESULTS


Taxonomic Composition

The 16S rRNA gene sequencing is a rapid and accurate identification method for bacterial isolates; however, it is not applicable for several genera and only provides resolution till the genus level and the presence of nucleotide variations in rRNA operons in a single genome. Under this category, species distribution under the classification level of phylum up to the classification level of the genus was done. The distribution histogram of the relative abundance was generated as shown in **Figures 2, 3**. The DADA2 pipeline of colon mucosa samples was classified into six bacterial phyla. Overall, *Firmicutes* (85.41%) in T2, *Bacteroidetes* (40.24%) in T3, and *Proteobacteria* (10.03%) in T7 were the three most dominant phyla (**Figure 2**). The overall genus showed that *Bacteroides* (39.37%) in T3, *Faecalibacterium* (17.35%) in T5, *Lactobacillus* (14.39%) in T6, *Ruminococcaceae* UCG-14 (11.01%) in T2, *Escherichia-Shigella* (10.33%) in T7, and (*Ruminococcus*) torques group (4.82%) in T2 were the dominant genera (**Figure 3**).

Species Diversity

Alpha Diversity Indices of Colon Mucosa Microbiota

The alpha diversity of colon mucosa microbiota of broiler chickens fed postbiotics and paraprobiotics was measured through five indices: Observed, Chao1, Shannon, Simpson, and Fisher (**Figure 4**). The Observed and Chao1 indices showed increases in richness in the paraprobiotics, positive control, and postbiotics groups. Paraprobiotic RG11 (T5) was recorded to have a higher richness mean value of above 160 ASVs according to the Chao1 index. Fisher's index showed higher






FIGURE 3 | Relative abundance in % of colon microbiota at the genus level. T1 = Negative control (Basal diet), T2 = Positive control [Basalt diet + 0.01% (w/w) Oxytetracycline], T3 = Basal diet + 0.2% (v/w) postbiotic TL1, T4 = Basal diet + 0.2% (v/w) Postbiotic RS5, T5 = Basal diet + 0.2%(v/w) paraprobiotic RG11, T6 = Basal diet + 0.2% (v/w) postbiotic RI11, T7 = Basal diet + 0.2% (v/w) paraprobiotic RG14, T8 = Basal diet + 0.2% (v/w) paraprobiotic RI11. *Bacteroides* (39.36%) in T3, *Faecalibacterium* (17.35%) in T5, *Lactobacillus* (14.39%) in T6, *Ruminococcaceae* UCG-14 (11.01%) in T2, *Escherichia-Shigella* (10.33%) in T7 are the most dominant genus.





FIGURE 4 | Alpha diversity indices of colon mucosa microbiota of broiler chickens fed postbiotics and paraprobiotics. T1 = Negative control (Basal diet), T2 = Positive control [Basalt diet + 0.01% (w/w) Oxytetracycline], T3 = Basal diet + 0.2% (v/w) postbiotic TL1, T4 = Basal diet + 0.2% (v/w) Postbiotic RS5, T5 = Basal diet + 0.2%(v/w) paraprobiotic RG11, T6 = Basal diet + 0.2% (v/w) postbiotic RI11, T7 = Basal diet + 0.2% (v/w) paraprobiotic RG14, T8 = Basal diet + 0.2% (v/w) paraprobiotic RI11.



species richness in the paraprobiotics, positive control, and postbiotics groups. According to Fisher's index, paraprobiotic RG11 (T5) had the highest richness mean value above 25 ASVs. Shannon's index showed more species diversity in T2 and T5 than the other treatment groups. The positive control (T2) and T5 were recorded to have higher species diversity of above 4.5 on Shannon's index. Similarly, the positive control (T2) and T5 had a higher value on Simpson's index at above 0.985.

The sample-based rarefaction curve in **Figure 5** matched the previously determined maximum species richness by Chao1 and Fisher of alpha diversity. Paraprobiotic RG11 (T5) still maintained the highest richness based on the rarefaction curve. According to the rarefaction curves, all the curves of the treatment group reached their plateau, indicating that the read depth was sufficient and less new (new species) can be detected with increasing sequencing depth.

Shared and Unique Microbial Composition

The comparison of the control and postbiotics groups showed that 38 ASVs were common to the five treatment groups, and 146 more unique ASVs were found in T6 (**Figure 6A**). On the other hand, the comparison between the control groups and paraprobiotic groups showed that a total of 45 ASVs were common to the five treatment groups, and 144 more unique ASVs were found in T5 (**Figure 6B**). The Venn diagram in **Figure 6C** represents the comparison of ASVs between the postbiotics and paraprobiotics groups. The results showed that

out of the total abundance ASVs found in the six treatments, T5 had 133 more unique ASVs, and 36 ASVs were common to both the postbiotics and paraprobiotics groups.


Phylogenetic Tree

The relationship between the bacterial species was studied by constructing a phylogenetic tree using a combination of MAFFT and FastTree algorithms based on maximum-likelihood NNIs. The phylogenetic tree confirms the genus to which the query sequence strain belongs and its closest neighbours by comparing it with other sequences from the database. Further, genotypic, chemotaxonomic, and phenotypic analysis platforms are designed. The overall most closely related genus among the 30 top genera according to ASVs in colon mucosa microbiota of the eight dietary treatments was the genus *Bacteroides* (see **Figure 7**).

DISCUSSION

Effect of Postbiotics and Paraprobiotics on 16S RRNA Amplicon Sequencing of Colon Mucosa Microbiota


According to (36), the GIT of chickens harbours a large microbial community that plays an important role in chicken growth and health through enhancing nutrient absorption and strengthening the immune system. The GIT provides a platform for the growth of a diverse microbiota that provides a second barrier against colonisation by pathogens, regulates immune development and



maturation, and provides metabolites for host nutrition (37–39). SCFA, such as butyrate, provides energy for the epithelial cells that line the intestine and subdues the expression of virulence factors of harmful (38, 40–42).

There is no doubt that caecal microbiota has the greatest diversity in the GI tract. It is the source for most, if not

all, mucosa-associated microbiota of the proximal and distal colon (43). Until now, most studies on chicken intestinal microbiota are focused more on the caecal section of the hindgut. However, using 16S rRNA amplicon sequence, we focused the current research on the colon mucosa microbiota of broiler chickens.



The six most abundant phyla reported in this study are Firmicutes, Bacteroidetes, Proteobacteria, Verrucomicrobia, Tenericutes, and Actinobacteria in the colon mucosa. Similarly, the five most abundant phyla reported in the duodenum, jejunum, ileum, colon, and caecum microbiota were Firmicutes, Bacteroidetes, Proteobacteria, Actinobacteria, and Cyanobacteria (44); this elucidated the fact that caecal microbiota was the major source of mucosa-associated microbiota of the distal colon (43). According to our findings, Firmicutes was the dominant phylum in all the treatment groups with a significantly higher abundance of 85.41% in T2 and 82.66% in T5 than other treatments. The alpha diversity indices (Chao1, Fisher, Observed, Shannon, and Simpson) indicated an increase in species diversity and richness in the colon mucosa microbiota.

The implication of the increase in abundance of Firmicutes as a result of dietary supplementation with postbiotics and paraprobiotics in this study is more butyrate production. Firmicutes were reported as the phylum with a larger number of taxa encoding enzymes required for butyrate production (45). Previously, postbiotic metabolite combination was reported to increase the faecal butyric acid concentrations in broiler chickens

(14). Furthermore, butyrate is the main source of energy for enterocytes, and it helps regulate cellular differentiation and proliferation within the intestinal mucosa, thereby increasing intestinal tissue weight (40, 46–48). The stimulation of the release of gastrointestinal peptides and growth factors by butyrate acting on cell proliferation is one of the major mechanisms involved in intestinal mucosa proliferation by butyrate (46). In addition, previous research findings revealed that butyrate increases the secretion of IL-10 and decreases the secretion of interferon- γ by activated human lymphocytes *in vitro* (49, 50). Also, there was a reported decrease in *ex vivo* production of inflammatory cytokines in intestinal biopsies of humans who have Crohn's disease and a reduction in the severity of 2,4,6-trinitrobenzene sulfonic acid-induced colitis in rats caused by butyrate (51).

Bacteroides (39.37%), *Faecalibacterium* (17.35%), *Lactobacillus* (14.39%), *Ruminococcaceae* UCG-014 (11.01%), and *Escherichia-Shigella* (10.33%) were the most dominant out of the overall genera sequenced in this study. Similarly, it was reported that *Streptococcus*, uncultured *Ruminococcaceae*, and *Lactobacillus* were the three most predominant genera in the colonic digesta and mucosa of pigs (52). The production of

propionate and succinate was associated with *Bacteroides* as terminal products of metabolism, as reported by Adamberg et al. (53). Propionate is a less preferred substrate of colonocytes but is transported to the liver and used as an important energy source for the host (47). The strain *Faecalibacterium* was reported to be a carrier of the enzymes necessary for butyrate production and present from the early stages of development. Therefore, the strain will actively participate in future intervention and modulation of the gut microbiota by improving the overall health and growth performance of poultry (45). It was revealed recently that some *Ruminococcus* species in the human colon were found to play a primary role in the degradation of dietary resistant starch (54). *Lactobacillus*, an important probiotic bacterium in promoting a healthy gut, was the fourth most predominant genus. A recent study with postbiotics also revealed a significant ($p < 0.05$) increase in the population of *Lactobacillus* in the caecum of broiler chickens (17). *Lactobacillus* is a beneficial microbe that can produce bacteriocins, a natural antimicrobial compound capable of inhibiting the growth of pathogens at molecular and cellular levels (55). The presence of *Lactobacillus* could explain why the population of chicken pathogens was significantly inhibited in this study. The phylum Proteobacteria in the colon mucosa was greatly decreased by postbiotics, paraprobiotics, and positive control, except for paraprobiotic RG14 (T7), where its population was higher.

CONCLUSION

This study revealed that supplementation of postbiotics and paraprobiotics in the broilers' chicken diet demonstrated a

positive effect on the microbiota by supporting the increase of beneficial microbes like the Firmicutes while decreasing harmful microbes like the Proteobacteria. The modification in the microbiota can result in a healthier gut. Therefore, postbiotics and paraprobiotics can positively affect the microbiota of the colon mucosa.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author/s.

ETHICS STATEMENT

The animal study was reviewed and approved by Institutional Animal Care and Use Committee (IACUC) with a Reference No: UPM/IACUC/AUP-R098/2018.

AUTHOR CONTRIBUTIONS

YD, TCL, and HLF designed the study. HLF and TCL provided probiotic strains and methods to produce postbiotics, probiotic suspension, and paraprobiotics and modified the manuscript. YD and AMN helped to perform the experiments. YD analysed the data. HA and NAMT assisted in proofreading the manuscript. All authors contributed to the article and approved the submitted version.

REFERENCES

- Shang Y, Kumar S, Oakley B, Kim WK. Chicken gut microbiota: importance and detection technology. *Front Vet Sci.* (2018) 5:1–11. doi: 10.3389/fvets.2018.00254
- Gaskins HR, Collier CT, Anderson DB. Antibiotics as growth promotants: mode of action. *Anim Biotechnol.* (2002) 13:29–42. doi: 10.1081/ABIO-120005768
- Jeurissen SH, Lewis M, van der Klis F, Mroz JD, Rebel Z, ter Huurne AAHM. Parameters and techniques to determine intestinal health of poultry as constituted by immunity, integrity, and functionality. *Curr Issues Intestinal Microbiol.* (2002) 3:1–14.
- Kohl KD. Diversity and function of the avian gut microbiota. *J Comp Physiol B: Biochem Syst Environ Physiol.* (2012) 182:591–602. doi: 10.1007/s00360-012-0645-z
- Wei S, Morrison M, Yu Z. Bacterial census of poultry intestinal microbiome. *Poult Sci.* (2013) 92:671–83. doi: 10.3382/ps.2012-02822
- Yegani M, Korver DR. Factors affecting intestinal health in poultry. *Poult Sci.* (2008) 87:2052–63. doi: 10.3382/ps.2008-00091
- Dibner JJ, Richards JD. Antibiotic growth promoters in agriculture: history and mode of action. *Poult Sci.* (2005) 84:634–43. doi: 10.1093/ps/84.4.634
- Shakouri MD, Iji PA, Mikkelsen LL, Cowieson AJ. Intestinal function and gut microflora of broiler chickens as influenced by cereal grains and microbial enzyme supplementation. *J Anim Physiol Anim Nutr.* (2009) 93:647–58. doi: 10.1111/j.1439-0396.2008.00852.x
- Oakley BB, Lillehoj HS, Kogut MH, Kim WK, Maurer JJ, Pedrosa A, et al. The chicken gastrointestinal microbiome. *FEMS Microbiol Lett.* (2014) 360:100–12. doi: 10.1111/1574-6968.12608
- Ricke SC. Perspectives on the use of organic acids and short chain fatty acids as antimicrobials. *Poult Sci.* (2003) 82:632–9. doi: 10.1093/ps/82.4.632
- Christl SU, Bartram HP, Paul A, Kelber E, Scheppach W, Kasper H. Bile acid metabolism by colonic bacteria in continuous culture: Effects of starch and pH. *Ann Nutr Metab.* (1997) 41:45–51. doi: 10.1159/000177977
- Pan D, Yu Z. Intestinal microbiome of poultry and its interaction with host and diet. *Gut Microbes.* (2014) 5:108–19. doi: 10.4161/gmic.26945
- Borda-Molina D, Seifert J, Camarinha-Silva A. Current perspectives of the chicken gastrointestinal tract and its microbiome. *Comput Struct Biotechnol J.* (2018) 16:131–9. doi: 10.1016/j.csbj.2018.03.002
- Thanh NT, Loh TC, Foo HL, Azhar BK. Effects of feeding metabolite combinations produced by *Lactobacillus plantarum* on growth performance, faecal microbial population, small intestine villus height and faecal volatile fatty acids in broilers. *Br Poult Sci.* (2009) 50:298–306. doi: 10.1080/00071660902873947
- Loh TC, Thanh NT, Foo HL, Hair-Bejo M, Azhar BK. Feeding of different levels of metabolite combinations produced by *Lactobacillus plantarum* on growth performance, fecal microflora, volatile fatty acids and villi height in broilers. *Anim Sci J.* (2010) 81(2), 205–214. doi: 10.1111/j.1740-0929.2009.00701.x
- Kareem KY, Loh TC, Foo HL, Asmara SA, Akit H. Influence of postbiotic RG14 and inulin combination on cecal microbiota, organic acid concentration, and cytokine expression in broiler chickens. *Poult Sci.* (2017) 96:966–75. doi: 10.3382/ps/pew362
- Humam AM, Loh TC, Foo HL, Samsudin AA, Mustapha NM, Zulkifli I, et al. Effects of feeding different postbiotics produced by *Lactobacillus plantarum* on growth performance, carcass yield, intestinal morphology, gut microbiota composition, immune status, and growth gene expression in broilers under heat stress. *Animals.* (2019) 9:1–20. doi: 10.3390/ani9090644

18. Tsilingiri K, Barbosa T, Penna G, Caprioli F, Sonzogni A, Viale G, et al. Probiotic and postbiotic activity in health and disease: comparison on a novel polarised ex-vivo organ culture model. *Gut*. (2012) 61:1007–15. doi: 10.1136/gutjnl-2011-300971
19. Tsilingiri K, Rescigno M. Postbiotics: what else? *Benef Microbes*. (2013) 4:101–7. doi: 10.3920/BM2012.0046
20. Taverniti V, Guglielmetti S. The immunomodulatory properties of probiotic microorganisms beyond their viability (ghost probiotics: proposal of paraprobiotic concept). *Genes Nutr*. (2011) 6:261–74. doi: 10.1007/s12263-011-0218-x
21. Thanh NT, Chwen LT, Foo HL, Hair-bejo M, Kasim AB. Inhibitory activity of metabolites produced by strains of *Lactobacillus plantarum* isolated from Malaysian fermented food. *Int J Probiotics Prebiotics*. (2010) 5:37–44.
22. Fujiki T, Hirose Y, Yamamoto Y, Murosaki S. Enhanced immunomodulatory activity and stability in simulated digestive juices of *Lactobacillus plantarum* L-137 by heat treatment. *Biosci Biotechnol Biochem*. (2012) 76:918–22. doi: 10.1271/bbb.110919
23. Kareem KY, Ling FH, Chwen LT, Foong OM, Asmara SA. Inhibitory activity of postbiotic produced by strains of *Lactobacillus plantarum* using reconstituted media supplemented with inulin. *Gut Pathog*. (2014) 6:23–9. doi: 10.1186/1757-4749-6-23
24. Ou CC, Lin SL, Tsai JJ, Lin MY. Heat-killed lactic acid bacteria enhance immunomodulatory potential by skewing the immune response toward Th1 polarization. *J Food Sci*. (2011) 76:M260–7. doi: 10.1111/j.1750-3841.2011.02161.x
25. Villena J, Barbieri N, Salva S, Herrera M, Alvarez S. Enhanced immune response to pneumococcal infection in mal-nourished mice nasally treated with heat-killed *Lactobacillus casei*. *Microbiol Immunol*. (2009) 53:636–46. doi: 10.1111/j.1348-0421.2009.00171.x
26. Loh TC, Choe DW, Foo HL, Sazili AQ, Bejo MH. Effects of feeding different postbiotic metabolite combinations produced by *Lactobacillus plantarum* strains on egg quality and production performance, faecal parameters and plasma cholesterol in laying hens. *BMC Vet Res*. (2014) 10:1–9. doi: 10.1186/1746-6148-10-149
27. Kareem, KY, Loh TC, Foo HL, Akit H, Samsudin AA. Effects of dietary postbiotic and inulin on growth performance, IGF1 and GHR mRNA expression, faecal microbiota and volatile fatty acids in broilers. *BMC Vet Res*. (2016) 12:8–17. doi: 10.1186/s12917-016-0790-9
28. Tlaskalová-Hogenová H, Třápková R, Kozáková H, Hudcovic T, Vannucci L, Tucková L, et al. The role of gut microbiota (commensal bacteria) and the mucosal barrier in the pathogenesis of inflammatory and autoimmune diseases and cancer: Contribution of germ-free and gnotobiotic animal models of human diseases. *Cell Mol Immunol*. (2011) 8:110–20. doi: 10.1038/cmi.2010.67
29. Burrough ER, Arruda BL, Plummer PJ. Comparison of the luminal and mucosa-associated microbiota in the colon of pigs with and without swine dysentery. *Front Vet Sci*. (2017) 4:1–10. doi: 10.3389/fvets.2017.00139
30. Mohamad ZNA, Foo HL, Loh TC, Mohamad R, Rahim RA. Enhancement of versatile extracellular cellulolytic and hemicellulolytic enzyme productions by *Lactobacillus plantarum* RI 11 isolated from Malaysian food using renewable natural polymers. *Molecules*. (2020) 25:1–7. doi: 10.3390/molecules25112607
31. Ooi MF, Nurzafrah M, Hooi LF, Teck CL, Rosfarizan M, Raha AR, et al. Effects of carbon and nitrogen sources on bacteriocin- inhibitory activity of postbiotic metabolites produced by *Lactobacillus plantarum* I-UL4. *Malays J Microbiol*. (2015) 11:176–84. doi: 10.1017/CBO9781107415324.004
32. Callahan B. *DADA2 Pipeline Tutorial (1.16)* (2020). Available online at: <https://benjineb.github.io/dada2/tutorial.html>
33. Xue Z, Kable ME, Marco ML. Impact of DNA sequencing and analysis methods on 16S rRNA gene bacterial community analysis of dairy products. *MSphere*. (2018) 3:1–14. doi: 10.1128/msphere.00410-18
34. Nearing JT, Douglas GM, Comeau AM, Langille MGI. Denoising the Denoisers: an independent evaluation of microbiome sequence error—correction approaches. *PeerJ*. (2018) 2018:1–22. doi: 10.7717/peerj.5364
35. Callahan BJ, McMurdie PJ, Rosen, MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High resolution sample inference from Illumina amplicon data. *Nat Methods*. (2016) 13:581–3. doi: 10.1038/nmeth.3869
36. Choi KY, Lee TK, Sul WJ. Metagenomic analysis of chicken gut microbiota for improving metabolism and health of chickens—a review. *Asian Aust J Anim Sci*. (2015) 28:1217–25. doi: 10.5713/ajas.15.0026
37. Gaggia F, Mattarelli P, Biavati B. Probiotics and prebiotics in animal feeding for safe food production. *Int J Food Microbiol*. (2010) 141(SUPPL.): S15–28. doi: 10.1016/j.ijfoodmicro.2010.02.031
38. Sergeant MJ, Constantinidou C, Cogan TA, Bedford MR, Penn CW, Pallen MJ. Extensive microbial and functional diversity within the chicken cecal microbiome. *PLoS ONE*. (2014) 9:1–13. doi: 10.1371/journal.pone.0091941
39. Roberts T, Wilson J, Guthrie A, Cookson K, Vancraeynest D, Schaeffer J, et al. New issues and science in broiler chicken intestinal health: Emerging technology and alternative interventions. *J Appl Poultry Res*. (2015) 24:257–66. doi: 10.3382/japr/pfv023
40. Rinttilä T, Apajalahti J. Intestinal microbiota and metabolites—Implications for broiler chicken health and performance. *J Appl Poultry Res*. (2013) 22:647–58. doi: 10.3382/japr.2013-00742
41. Roto SM, Rubinelli PM, Ricke SC. An introduction to the avian gut microbiota and the effects of yeast-based prebiotic-type compounds as potential feed additives. *Front Vet Sci*. (2015) 2:1–27. doi: 10.3389/fvets.2015.00028
42. Polansky O, Sekelova Z, Faldynova M, Sebkova A, Sisak F, Rychlik I. Important metabolic pathways and biological processes expressed by chicken cecal microbiota. *Appl Environ Microbiol*. (2016) 82:1569–76. doi: 10.1128/AEM.03473-15
43. Wang Y, Devkota S, Musch MW, Jabri B, Nagler C, Antonopoulos DA, et al. Regional mucosa-associated microbiota determines physiological expression of TLR2 and TLR4 in murine colon. *PLoS ONE*. (2010) 5:1–12. doi: 10.1371/journal.pone.0013607
44. Xiao Y, Xiang Y, Zhou W, Chen J, Li K, Yang H. Microbial community mapping in intestinal tract of broiler chicken. *Poult Sci*. (2017) 96:1387–93. doi: 10.3382/ps/pew372
45. Segura-Wang M, Grabner N, Koestelbauer A, Klose V, Ghanbari M. Genome-resolved metagenomics of the chicken gut microbiome. *Front Microbiol*. (2021) 12:1–13. doi: 10.3389/fmicb.2021.726923
46. Guilloteau P, Martin L, Eeckhaut V, Ducatelle R, Zabielski R, Van Immerseel F. From the gut to the peripheral tissues: the multiple effects of butyrate. *Nutr Res Rev*. (2010) 23:366–84. doi: 10.1017/S0954422410000247
47. Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F. From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell*. (2016) 165:1332–45. doi: 10.1016/j.cell.2016.05.041
48. Beauclercq S, Nadal-Desbarats L, Hennequet-Antier C, Gabriel I, Tesseraud S, Calenge F, et al. Relationships between digestive efficiency and metabolomic profiles of serum and intestinal contents in chickens. *Sci Rep*. (2018) 8:1–11. doi: 10.1038/s41598-018-24978-9
49. Säemann MD, Böhmig GA, Österreicher CH, Burtscher H, Parolini O, Diakos C, et al. Anti-inflammatory effects of sodium butyrate on human monocytes: potent inhibition of IL-12 and up-regulation of IL-10 production. *FASEB J*. (2000) 14:2380–2. doi: 10.1096/fj.00-0359fj
50. Nancey S, Bienvenu J, Coffin B, Andre F, Descos L, Flourié B. Butyrate strongly inhibits in vitro stimulated release of cytokines in blood. *Dig Dis Sci*. (2002) 47:921–8. doi: 10.1023/A:1014781109498
51. Segain JP, Raingeard de la Blétière D, Bourreille A, Leray V, Gervois N, Rosales C, et al. Butyrate inhibits inflammatory responses through NFκB inhibition: implications for Crohn's disease. *Gut Pathogens*. (2000) 47:397–403. doi: 10.1136/gut.47.3.397
52. Sun Y, Zhou L, Fang L, Su Y, Zhu W. Responses in colonic microbial community and gene expression of pigs to a long-term high resistant starch diet. *Front Microbiol*. (2015) 6:1–10. doi: 10.3389/fmicb.2015.00877
53. Adamberg S, Tomson K, Vija H, Puurand M, Kabanova N, Visnapuu T, et al. Degradation of fructans and production of propionic acid by bacteroides thetaiotaomicron are enhanced by the shortage of amino acids. *Front Nutr*. (2014) 1:1–10. doi: 10.3389/fnut.2014.00021
54. Ze X, Duncan SH, Louis P, Flint HJ. Ruminococcus bromii is a keystone species for the degradation of resistant starch in the human colon. *ISME J*. (2012) 6:1535–43. doi: 10.1038/ismej.2012.4

55. Drider D, Fimland G, Héchard Y, McMullen LM, Prévost H. The continuing story of class IIa bacteriocins. *Microbiol Mol Biol Rev.* (2006) 70:564–82. doi: 10.1128/membr.00016-05

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of

the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Danladi, Loh, Foo, Akit, Md Tamrin and Mohammad Naeem. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.